

2017

Characterizing the Volatilome of Land-disposed Sewage Sludge Under Seasonal Temperature Regimes

Hope L. Juntunen
Dakota State University

Lucas Leinen
Dakota State University

Vaille Swenson
Dakota State University

R C. Hale
Dakota State University

R Honour
Dakota State University

See next page for additional authors

Follow this and additional works at: <https://scholar.dsu.edu/anspapers>

 Part of the [Environmental Health and Protection Commons](#), [Other Environmental Sciences Commons](#), and the [Other Physical Sciences and Mathematics Commons](#)

Recommended Citation

Juntunen, Hope L.; Leinen, Lucas; Swenson, Vaille; Hale, R C.; Honour, R; Gaylor, Michael; and Videau, Patrick, "Characterizing the Volatilome of Land-disposed Sewage Sludge Under Seasonal Temperature Regimes" (2017). *Faculty Research & Publications*. 7.
<https://scholar.dsu.edu/anspapers/7>

This Conference Proceeding is brought to you for free and open access by the College of Arts & Sciences at Beadle Scholar. It has been accepted for inclusion in Faculty Research & Publications by an authorized administrator of Beadle Scholar. For more information, please contact repository@dsu.edu.

Authors

Hope L. Juntunen, Lucas Leinen, Vaille Swenson, R C. Hale, R Honour, Michael Gaylor, and Patrick Videau

CHARACTERIZING THE VOLATILOME OF LAND-DISPOSED SEWAGE SLUDGE UNDER SEASONAL TEMPERATURE REGIMES

H. L. Juntunen¹, L. J. Leinen¹, V. A. Swenson¹, R. Honour², R. C. Hale³,
P. Videau¹, and M. O. Gaylor^{1*}

¹College of Arts and Science
Dakota State University
Madison, SD 50742

²The Precautionary Group LLC
Kenmore, WA 98028

³Virginia Institute of Marine Science
College of William and Mary
Gloucester Point, VA 23062

*Corresponding author email: michael.gaylor@dsu.edu

ABSTRACT

About eight million dry tons of sewage sludge waste is generated in the US annually, with more than half of that now land-disposed on agricultural and forested lands. Though containing essential plant nutrients, sludge also harbors complex mixtures of volatile organic compounds (VOCs) that result in toxic emissions therefrom. While ecotoxic impacts to sludged ecosystems are a primary concern, the stifling emissions are most obvious to and disconcerting for the public, which has led to increasing concerns for the safety of this practice. The large-scale disposal of sludge in the temperate rainforests of the Puget Sound Watershed has resulted in acute toxicity to macrobiota, and despite escalating concerns for detrimental impacts, little is known about the total VOC inventories, or “volatilomes,” of these complex wastes. To address this knowledge gap, we characterized VOC emissions from forest-disposed sewage sludge over a range of seasonal temperature regimes. We also incubated sludge samples at the more extreme 100 °C to assess the “complete volatilome.” After 1-hr incubations in gastight vials, VOCs accumulated in the headspace were sampled with a gastight syringe and analyzed with gas chromatography-mass spectrometry to generate distinct chemical fingerprints of sludge sample volatilomes over the range of temperatures. Total integrated chromatographic peak areas increased with temperature, indicating increased VOC production. Sludge volatilomes were dominated by a multitude of aliphatics and aromatics, with comparatively lesser emissions of alcohols, esters, aldehydes, terpenes, and nitrogen-, sulfur-, and halogen-containing compounds.